

# Eigenvalue asymptotics for the Schrödinger operator with a $\delta$ -interaction on a punctured surface

P. Exner<sup>a,b</sup> and K. Yoshitomi<sup>c</sup>

- a) Department of Theoretical Physics, Nuclear Physics Institute, Academy of Sciences, 25068 Řež, Czech Republic*
- b) Doppler Institute, Czech Technical University, Břehová 7, 11519 Prague, Czech Republic*
- c) Department of Mathematics, Tokyo Metropolitan University, Minami-Ohsawa 1-1, Hachioji-shi, Tokyo 192-0397, Japan  
exner@ujf.cas.cz, yositomi@comp.metro-u.ac.jp*

MSC numbers: 35J10, 81V99

KEYWORDS: Schrödinger operators, singular interaction, discrete spectrum, manifolds, perturbation

**Abstract.** Given  $n \geq 2$ , we put  $r = \min\{i \in \mathbb{N}; i > n/2\}$ . Let  $\Sigma$  be a compact,  $C^r$ -smooth surface in  $\mathbb{R}^n$  which contains the origin. Let further  $\{S_\epsilon\}_{0 \leq \epsilon < \eta}$  be a family of measurable subsets of  $\Sigma$  such that  $\sup_{x \in S_\epsilon} |x| = \mathcal{O}(\epsilon)$  as  $\epsilon \rightarrow 0$ . We derive an asymptotic expansion for the discrete spectrum of the Schrödinger operator  $-\Delta - \beta\delta(\cdot - \Sigma \setminus S_\epsilon)$  in  $L^2(\mathbb{R}^n)$ , where  $\beta$  is a positive constant, as  $\epsilon \rightarrow 0$ . An analogous result is given also for geometrically induced bound states due to a  $\delta$  interaction supported by an infinite planar curve.

# 1 Introduction

Schrödinger operators with  $\delta$ -interactions supported by subsets of a lower dimension in the configuration space have been studied by numerous authors – see, e.g., [2]–[5] and references therein. Recently such systems attracted a new attention as models of “leaky” quantum wires and similar structures; new results have been derived about a curvature-induced discrete spectrum [7, 8] and the strong-coupling asymptotics [6, 9, 10, 11, 12].

The purpose of this paper is to discuss another question, namely how the discrete spectra of such operators behave with respect to a perturbation of the interaction support. Since the argument we are going to use can be formulated in any dimension, we consider here generally  $n$ -dimensional Schrödinger operators,  $n \geq 2$ , with a  $\delta$ -interaction supported by a punctured surface. On the other hand, we restrict our attention to the situation when the surface codimension is one and the Schrödinger operator in question is defined naturally by means of the appropriate quadratic form.

Formally speaking, our result says that up to an error term the eigenvalue shift resulting from removing an  $\epsilon$ -neighbourhood of a surface point is the same as that of adding a repulsive  $\delta$  interaction at this point with the coupling constant proportional to the puncture “area”. We will formulate this claim precisely in Theorem 1 below for any sufficiently smooth compact surface in  $\mathbb{R}^n$  and prove it in Section 3. Furthermore, the compactness requirement is not essential in the argument; in Section 4 we will derive an analogous asymptotic formula for an infinite planar curve which is not a straight line but it is asymptotically straight in a suitable sense.

# 2 The main result

Put  $r := \min\{i \in \mathbb{N}; i > n/2\}$ . Let  $\Sigma$  be a compact,  $C^r$ -smooth surface in  $\mathbb{R}^n$  which contains the origin,  $0 \in \Sigma$ . Let further  $\{S_\epsilon\}_{0 \leq \epsilon < \eta}$  be a family of subsets of  $\Sigma$  which obeys the following hypotheses:

- (H.1) The set  $S_\epsilon$  is measurable with respect to the  $(n-1)$ -dimensional Lebesgue measure on  $\Sigma$  for any  $\epsilon \in [0, \eta)$ .
- (H.2)  $\sup_{x \in S_\epsilon} |x| = \mathcal{O}(\epsilon)$  as  $\epsilon \rightarrow 0$ .

Next we fix  $\beta > 0$  and define for  $0 \leq \epsilon < \eta$  the quadratic form  $q_\epsilon$  by

$$q_\epsilon[u, v] := (\nabla u, \nabla v)_{L^2(\mathbb{R}^n)} - \beta \int_{\Sigma \setminus S_\epsilon} u(x) \overline{v(x)} dS, \quad u, v \in H^1(\mathbb{R}^n);$$

it is easily seen to be closed and bounded from below. Let  $H_\epsilon$  be the self-adjoint operator associated with  $q_\epsilon$ . Since  $\Sigma \setminus S_\epsilon$  is bounded, we have

$$\sigma_{\text{ess}}(H_\epsilon) = [0, \infty) \quad \text{and} \quad \sharp \sigma_{\text{disc}}(H_0) < \infty.$$

By the min-max principle, there exists a unique  $\beta^* \geq 0$  such that  $\sigma_{\text{disc}}(H_0)$  is non-empty if  $\beta > \beta^*$  while  $\sigma_{\text{disc}}(H_0) = \emptyset$  for  $\beta \leq \beta^*$ . The critical coupling is dimension-dependent: a straightforward modification of the usual Birman-Schwinger argument using [5, Lemma 2.3] shows that  $\beta^* = 0$  when  $n = 2$ , while for  $n \geq 3$  we have  $\beta^* > 0$  by [5, Thm 4.2(iii)]. Since our aim is to derive asymptotic properties of the discrete spectrum, we will assume throughout that

$$(H.3) \quad \beta > \beta^*.$$

Let  $N$  be the number of negative eigenvalues of  $H_0$ . Since

$$0 \leq q_\epsilon[u, u] - q_0[u, u] \rightarrow 0 \quad \text{as} \quad \epsilon \rightarrow 0 \quad \text{for} \quad u \in H^1(\mathbb{R}^n),$$

there exists  $\eta' \in (0, \eta)$  such that for  $\epsilon \in (0, \eta')$  the operator  $H_\epsilon$  has exactly  $N$  negative eigenvalues denoted by  $\lambda_1(\epsilon) < \lambda_2(\epsilon) \leq \dots \leq \lambda_N(\epsilon)$ , and moreover

$$\lambda_j(\epsilon) \rightarrow \lambda_j(0) \quad \text{as} \quad \epsilon \rightarrow 0 \quad \text{for} \quad 1 \leq j \leq N$$

(see [14, Chap. VIII, Thm 3.15]). Let  $\{\varphi_j(x)\}_{j=1}^N$  be an orthonormal system of eigenfunctions of  $H_0$  such that  $H_0 \varphi_j = \lambda_j(0) \varphi_j$  for  $1 \leq j \leq N$ . Pick a sufficiently small  $a > 0$  so that the set  $\{x \in \mathbb{R}^n : |x| < a\} \setminus \Sigma$  consists of two connected components, which we denote by  $B_\pm$ . We have  $\varphi_j \in H^r(B_\pm)$  by the elliptic regularity theorem (see [1, Sec. 10]), because the form domain  $H^1(\mathbb{R}^n)$  of  $q_0$  is locally invariant under tangential translations along the surface  $\Sigma$ . Since  $r > n/2$  by assumption, the Sobolev trace theorem implies that the function  $\varphi_j$  is continuous on a  $\Sigma$ -neighbourhood of the origin. We also note that one can suppose without loss of generality that  $\varphi_1(x) > 0$  in  $\mathbb{R}^n$ . For a given  $\mu \in \sigma_{\text{disc}}(H_0)$  we define

$$\begin{aligned} m(\mu) &:= \min\{1 \leq j \leq N; \mu = \lambda_j(0)\}, \\ n(\mu) &:= \max\{1 \leq j \leq N; \mu = \lambda_j(0)\}, \\ C(\mu) &:= \left( \varphi_i(0) \overline{\varphi_j(0)} \right)_{m(\mu) \leq i, j \leq n(\mu)}. \end{aligned}$$

Let  $s_{m(\mu)} \leq s_{m(\mu)+1} \leq \dots \leq s_{n(\mu)}$  be the eigenvalues of the matrix  $C(\mu)$ . In particular, if  $\mu = \lambda_j(0)$  is a simple eigenvalue of  $H_0$ , we have  $m(\mu) = n(\mu) = j$  and  $s_j = |\varphi_j(0)|^2$ . Our main result can be then stated as follows.

**Theorem 1** *Adopt the assumptions (H.1)–(H.3). Let  $\mu \in \sigma_{\text{disc}}(H_0)$ , then the asymptotic formula*

$$\lambda_j(\epsilon) = \mu + \beta \text{meas}_{\Sigma}(S_{\epsilon})s_j + o(\epsilon^{n-1}) \quad \text{as } \epsilon \rightarrow 0$$

*holds for  $m(\mu) \leq j \leq n(\mu)$ , where  $\text{meas}_{\Sigma}(\cdot)$  stands for the  $(n-1)$ -dimensional Lebesgue measure on  $\Sigma$ .*

It should be stressed that our problem involves a singular perturbation and thus it cannot be reduced to the general asymptotic perturbation theory of quadratic forms described in [14, Sec. VIII.4]. Indeed, we have

$$q_{\epsilon}[u, u] = q_0[u, u] + \beta \text{meas}_{\Sigma}(S_{\epsilon})|u(0)|^2 + \mathcal{O}(\epsilon^n) \quad \text{as } \epsilon \rightarrow 0$$

for  $u \in C_0^{\infty}(\mathbb{R}^n)$  and the quadratic form  $C_0^{\infty}(\mathbb{R}^n) \ni u \mapsto |u(0)|^2 \in \mathbb{R}$  does not extend to a bounded form on  $H^1(\mathbb{R}^n)$ , because the set

$$\{u \in C_0^{\infty}(\mathbb{R}^n); u = 0 \text{ in a neighbourhood of the origin}\}$$

is dense in  $H^1(\mathbb{R}^n)$ . We eliminate this difficulty by using the compactness of the map  $H^1(\mathbb{R}^n) \ni f \mapsto f|_{\Sigma} \in L^2(\Sigma)$ , which will enable us to prove Theorem 1 along the lines of the asymptotic-perturbation theorem proof.

Let us remark that our functional-analytic argument has a distinctive advantage over another technique employed in such situations, usually called the matching of asymptotic expansions – see [13] for a thorough review – since the latter typically requires a sort of self-similarity for the perturbation domains. Our technique needs no assumption of this type.

### 3 Proof of Theorem 1

We denote  $R(\zeta, \epsilon) = (H_{\epsilon} - \zeta)^{-1}$  for  $\zeta \in \rho(H_{\epsilon})$  and  $R(\zeta) = (H_0 - \zeta)^{-1}$  for  $\zeta \in \rho(H_0)$ . Put  $\kappa := \frac{1}{2} \text{dist}(\{\mu\}, \sigma(H_0) \setminus \{\mu\})$ . Since  $\lambda_j(\cdot)$  is continuous at the origin for  $1 \leq j \leq N$ , there is an  $\eta_0 \in (0, \eta')$  such that

$$\begin{aligned} \sigma(H_{\epsilon}) \cap [\mu - \kappa, \mu + \kappa] &= \sigma(H_{\epsilon}) \cap (\mu - \kappa/2, \mu + \kappa/2) \\ &= \{\lambda_{m(\mu)}(\epsilon), \lambda_{m(\mu)+1}(\epsilon), \dots, \lambda_{n(\mu)}(\epsilon)\} \end{aligned}$$

holds if  $0 < \epsilon \leq \eta_0$ . Choosing the circle  $C := \{z \in \mathbb{C}; |z - \mu| = \frac{3}{4}\kappa\}$  we put

$$w_j(\zeta, \epsilon) := R(\zeta, \epsilon)\varphi_j - R(\zeta)\varphi_j \quad \text{for } 0 < \epsilon \leq \eta_0, \zeta \in C.$$

Our first aim is to check that

$$\|w_j(\zeta, \epsilon)\|_{H^1(\mathbb{R}^n)} = \mathcal{O}(\epsilon^{(n-1)/2}) \quad \text{as } \epsilon \rightarrow 0 \quad (1)$$

holds uniformly with respect to  $\zeta \in C$  for  $m(\mu) \leq j \leq n(\mu)$ . Notice that there exists a  $K_0 > 0$  such that

$$\|u\|_{H^1(\mathbb{R}^n)}^2 \leq 2|(q_\epsilon - \zeta)[u, u]| + K_0\|u\|_{L^2(\mathbb{R}^n)}^2$$

for  $\zeta \in C$ ,  $u \in H^1(\mathbb{R}^n)$ , and  $0 < \epsilon \leq \eta_0$ . This implies that there exists a  $K_1 > 0$  such that

$$\|R(\zeta, \epsilon)u\|_{H^1(\mathbb{R}^n)} \leq K_1\|u\|_{L^2(\mathbb{R}^n)}$$

for  $\zeta \in C$ ,  $u \in H^1(\mathbb{R}^n)$ , and  $0 < \epsilon \leq \eta_0$ . Moreover, by the Sobolev trace theorem, there exists a constant  $K_2 > 0$  such that

$$\|u\|_{L^2(\Sigma)} \leq K_2\|u\|_{H^1(\mathbb{R}^n)} \quad \text{for } u \in H^1(\mathbb{R}^n).$$

Combining these three estimates we get

$$\begin{aligned} & \|w_j(\zeta, \epsilon)\|_{H^1(\mathbb{R}^n)}^2 \\ & \leq 2|(q_\epsilon - \zeta)[w_j(\zeta, \epsilon), w_j(\zeta, \epsilon)]| + K_0(w_j(\zeta, \epsilon), w_j(\zeta, \epsilon))_{L^2(\mathbb{R}^n)} \\ & = 2\left|-\beta \int_{S_\epsilon} R(\zeta)\varphi_j \overline{w_j(\zeta, \epsilon)} dS\right| + K_0(q_0 - q_\epsilon)[R(\zeta)\varphi_j, R(\bar{\zeta}, \epsilon)w_j(\zeta, \epsilon)] \\ & \leq \beta\|R(\zeta)\varphi_j\|_{L^2(S_\epsilon)}(2\|w_j(\zeta, \epsilon)\|_{L^2(S_\epsilon)} + K_0\|R(\bar{\zeta}, \epsilon)w_j(\zeta, \epsilon)\|_{L^2(S_\epsilon)}) \\ & = \frac{4\beta}{3\kappa}\|\varphi_j\|_{L^2(S_\epsilon)}(2\|w_j(\zeta, \epsilon)\|_{L^2(S_\epsilon)} + K_0\|R(\bar{\zeta}, \epsilon)w_j(\zeta, \epsilon)\|_{L^2(S_\epsilon)}) \\ & \leq \frac{4\beta}{3\kappa}K_2(2 + K_0K_1)\|\varphi_j\|_{L^2(S_\epsilon)}\|w_j(\zeta, \epsilon)\|_{H^1(\mathbb{R}^n)}. \end{aligned} \quad (2)$$

Since  $\|\varphi_j\|_{L^2(S_\epsilon)} = \mathcal{O}(\epsilon^{(n-1)/2})$  as  $\epsilon \rightarrow 0$  by the assumptions (H.1), (H.2) and the continuity of  $\varphi_j|_\Sigma$  at the origin, we arrive at the relation (1).

In the next step we are going to demonstrate that the convergence is in fact slightly faster, namely

$$\sup_{\zeta \in C} \|w_j(\zeta, \epsilon)\|_{H^1(\mathbb{R}^n)} = o(\epsilon^{(n-1)/2}) \quad \text{as } \epsilon \rightarrow 0. \quad (3)$$

We will proceed by contradiction. Suppose that (3) does not hold; then there would exist a constant  $\delta > 0$ , a sequence  $\{\epsilon_i\}_{i=1}^\infty \subset (0, \eta_0)$  which tends to zero, and  $\{\zeta_i\}_{i=1}^\infty \subset C$  such that

$$\epsilon_i^{-(n-1)/2} \|w_j(\zeta_i, \epsilon_i)\|_{H^1(\mathbb{R}^n)} \geq \delta \quad \text{for all } i \in \mathbb{N}. \quad (4)$$

Notice that the map  $H^1(\mathbb{R}^n) \ni f \mapsto f|_\Sigma \in L^2(\Sigma)$  is compact due to the boundedness of the map  $H^1(\mathbb{R}^n) \ni g \mapsto g|_\Sigma \in H^{1/2}(\Sigma)$  and the compactness of the imbedding  $H^{1/2}(\Sigma) \ni h \mapsto h \in L^2(\Sigma)$  – cf. [15, Chap. 1, Thms 8.3 and 16.1]. Since the two sequences

$$\left\{ \epsilon_i^{-(n-1)/2} w_j(\zeta_i, \epsilon_i) \right\}_{i=1}^\infty \quad \text{and} \quad \left\{ \epsilon_i^{-(n-1)/2} R(\overline{\zeta_i}, \epsilon_i) w_j(\zeta_i, \epsilon_i) \right\}_{i=1}^\infty$$

are bounded in  $H^1(\mathbb{R}^n)$ , there is a subsequence  $\{i(k)\}_{k=1}^\infty$  of  $\{i\}_{i=1}^\infty$  such that

$$\left\{ \epsilon_{i(k)}^{-(n-1)/2} w_j(\zeta_{i(k)}, \epsilon_{i(k)}) \right\}_{k=1}^\infty \quad \text{and} \quad \left\{ \epsilon_{i(k)}^{-(n-1)/2} R(\overline{\zeta_{i(k)}}, \epsilon_{i(k)}) w_j(\zeta_{i(k)}, \epsilon_{i(k)}) \right\}_{k=1}^\infty$$

converge in  $L^2(\Sigma)$ . Let us denote

$$g := \lim_{k \rightarrow \infty} \epsilon_{i(k)}^{-(n-1)/2} w_j(\zeta_{i(k)}, \epsilon_{i(k)}) \in L^2(\Sigma);$$

then we have

$$\begin{aligned} & \left\| \epsilon_{i(k)}^{-(n-1)/2} w_j(\zeta_{i(k)}, \epsilon_{i(k)}) \right\|_{L^2(S_{\epsilon_{i(k)}})} \\ & \leq \left\| \epsilon_{i(k)}^{-(n-1)/2} w_j(\zeta_{i(k)}, \epsilon_{i(k)}) - g \right\|_{L^2(\Sigma)} + \left( \int_{S_{\epsilon_{i(k)}}} |g(x)|^2 dS \right)^{1/2} \rightarrow 0 \end{aligned}$$

as  $k \rightarrow \infty$ . Similarly we obtain

$$\left\| \epsilon_{i(k)}^{-(n-1)/2} R(\overline{\zeta_{i(k)}}, \epsilon_{i(k)}) w_j(\zeta_{i(k)}, \epsilon_{i(k)}) \right\|_{L^2(S_{\epsilon_{i(k)}})} \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

Combining these result with the inequalities (2) we infer that

$$\epsilon_{i(k)}^{-(n-1)/2} \|w_j(\zeta_{i(k)}, \epsilon_{i(k)})\|_{H^1(\mathbb{R}^n)} \rightarrow 0 \quad \text{as } k \rightarrow \infty,$$

which violates the relation (4); in this way we have proved (3).

Now we denote by  $P_\epsilon$  the spectral projection of  $H_\epsilon$  associated with the interval  $(\mu - 3\kappa/4, \mu + 3\kappa/4)$ . It follows from (3) that

$$\begin{aligned} P_\epsilon \varphi_j - \varphi_j &= \frac{\sqrt{-1}}{2\pi} \oint_{|\zeta - \mu| = 3\kappa/4} w_j(\zeta, \epsilon) d\zeta \\ &= o(\epsilon^{(n-1)/2}) \quad \text{in } H^1(\mathbb{R}^n) \quad \text{as } \epsilon \rightarrow 0 \end{aligned}$$

holds for  $m(\mu) \leq j \leq n(\mu)$ . Consequently, we have

$$\begin{aligned} &(H_\epsilon P_\epsilon \varphi_i, P_\epsilon \varphi_j)_{L^2(\mathbb{R}^n)} - \mu \delta_{i,j} - \beta \varphi_i(0) \overline{\varphi_j(0)} \text{meas}_\Sigma(S_\epsilon) \\ &= q_\epsilon[P_\epsilon \varphi_i, P_\epsilon \varphi_j] - q_0[\varphi_i, \varphi_j] - \beta \varphi_i(0) \overline{\varphi_j(0)} \text{meas}_\Sigma(S_\epsilon) \\ &= q_\epsilon[\varphi_i, \varphi_j] - q_0[\varphi_i, \varphi_j] - q_\epsilon[(I - P_\epsilon)\varphi_i, (I - P_\epsilon)\varphi_j] \\ &\quad - \beta \varphi_i(0) \overline{\varphi_j(0)} \text{meas}_\Sigma(S_\epsilon) \\ &= -q_\epsilon[(I - P_\epsilon)\varphi_i, (I - P_\epsilon)\varphi_j] + \beta \int_{S_\epsilon} \varphi_i(x) \overline{\varphi_j(x)} dS \\ &\quad - \beta \varphi_i(0) \overline{\varphi_j(0)} \text{meas}_\Sigma(S_\epsilon) \\ &= o(\epsilon^{n-1}) \end{aligned} \tag{5}$$

and

$$(P_\epsilon \varphi_i, P_\epsilon \varphi_j)_{L^2(\mathbb{R}^n)} = \delta_{i,j} + o(\epsilon^{n-1}) \tag{6}$$

as  $\epsilon \rightarrow 0$  for  $m(\mu) \leq i, j \leq n(\mu)$ , where we have used, in the last step of (5), the assumptions (H.1), (H.2), the continuity of the restrictions  $\varphi_i|_\Sigma$  and  $\varphi_j|_\Sigma$  at the origin, and the uniform boundedness of  $q_\epsilon$  on  $H^1(\mathbb{R}^n)$  with respect to  $0 < \epsilon \leq \eta_0$ . Let us now introduce the matrices

$$\begin{aligned} L(\epsilon) &:= ((H_\epsilon P_\epsilon \varphi_i, P_\epsilon \varphi_j)_{L^2(\mathbb{R}^n)})_{m(\mu) \leq i, j \leq n(\mu)}, \\ M(\epsilon) &:= ((P_\epsilon \varphi_i, P_\epsilon \varphi_j)_{L^2(\mathbb{R}^n)})_{m(\mu) \leq i, j \leq n(\mu)}. \end{aligned}$$

Since  $\{P_\epsilon \varphi_j\}_{m(\mu) \leq j \leq n(\mu)}$  is a basis of the spectral subspace  $\text{Ran } P_\epsilon$ , we see that  $\lambda_{m(\mu)}(\epsilon), \lambda_{m(\mu)+1}(\epsilon), \dots, \lambda_{n(\mu)}(\epsilon)$  are the eigenvalues of the matrix  $L(\epsilon)M(\epsilon)^{-1}$ , which by (5), (6) is equal to

$$L(\epsilon)M(\epsilon)^{-1} = \mu I + \beta \text{meas}_\Sigma(S_\epsilon) C(\mu) + o(\epsilon^{n-1}),$$

where  $I$  stands for the identity matrix. This concludes the argument.  $\blacksquare$

## 4 Perturbation of an infinite curve

As we have mentioned, the compactness of  $\Sigma$  did not play an essential role in the above argument, and we can use the same technique for punctured non-compact manifolds of unit codimension as well, as long as the corresponding Hamiltonian has a discrete spectrum. At present this is known to be true in the case  $n = 2$  without restriction to the coupling constant  $\beta$ , see [7], and for  $n = 3$  and  $\beta$  large enough [9].

We shall thus consider “puncture” perturbations of infinite asymptotically straight curves. Let  $\Lambda : \mathbb{R} \rightarrow \mathbb{R}^2$  be a  $C^2$ -smooth curve parameterized by its arc length. Fix  $\beta > 0$  and assume that  $\Lambda(0) = 0$ . Given  $\epsilon \geq 0$ , we define

$$t_\epsilon[u, v] := (\nabla u, \nabla v)_{L^2(\mathbb{R}^2)} - \beta \int_{\Lambda(\mathbb{R} \setminus (-\epsilon, \epsilon))} u(x) \overline{v(x)} dS, \quad u, v \in H^1(\mathbb{R}^2).$$

Let  $T_\epsilon$  be the self-adjoint operator associated with the quadratic form  $t_\epsilon$ . We adopt the following assumptions about the curve  $\Lambda$ .

(H.4) The curve  $\Lambda$  is not a straight line.

(H.5) There exists  $c \in (0, 1)$  such that  $|\Lambda(s) - \Lambda(t)| \geq c|t - s|$  for  $s, t \in \mathbb{R}$ .

(H.6) There exist  $d > 0$ ,  $\rho > 1/2$ , and  $w \in (0, 1)$  such that the inequality

$$1 - \frac{|\Lambda(s) - \Lambda(s')|}{|s - s'|} \leq d [1 + |s + s'|^{2\rho}]^{-1/2}$$

holds in the sector  $\{(s, s') \in \mathbb{R}^2; w < \frac{s}{s'} < w^{-1}\}$ .

From [7, Prop 5.1 and Thm 5.2] we know that under these conditions

$$\sigma_{\text{ess}}(T_0) = [-\beta^2/4, \infty) \quad \text{and} \quad 1 \leq \#\sigma_{\text{disc}}(T_0) \leq \infty.$$

Let  $K := \{j \in \mathbb{N}; j \leq \#\sigma_{\text{disc}}(T_0)\}$ . For  $j \in K$ , we denote by  $\kappa_j(\epsilon)$  the  $j$ -th eigenvalue of  $T_\epsilon$  counted with multiplicity. The function  $\kappa_j(\cdot)$  is monotone non-decreasing, continuous function in a neighbourhood of the origin. Let  $\{\psi_j(x)\}_{j \in K}$  be an orthonormal system of eigenfunctions of  $T_0$  such that  $T_0 \psi_j = \kappa_j(0) \psi_j$  for  $j \in K$ . Each function  $\psi_j$  is continuous on  $\Lambda$ . For  $\mu \in \sigma_{\text{disc}}(T_0)$ , we define

$$\begin{aligned} p(\mu) &:= \min \{j \in K; \mu = \kappa_j(0)\}, \\ r(\mu) &:= \max \{j \in K; \mu = \kappa_j(0)\}, \\ D(\mu) &:= \left( \psi_i(0) \overline{\psi_j(0)} \right)_{p(\mu) \leq i, j \leq r(\mu)}. \end{aligned}$$



Let  $e_{p(\mu)} \leq e_{p(\mu)+1} \leq \dots \leq e_{r(\mu)}$  be the eigenvalues of the matrix  $D(\mu)$ . As in the compact case, if  $\mu = \kappa_j(0)$  is a simple eigenvalue of  $H_0$ , we have  $p(\mu) = r(\mu) = j$  and  $e_j = |\psi_j(0)|^2$ . The asymptotic behaviour now looks as follows.

**Theorem 2** *Assume that (H.4)–(H.6) and take  $\mu \in \sigma_{\text{disc}}(T_0)$ . Then*

$$\kappa_j(\epsilon) = \mu + 2\beta e_j \epsilon + o(\epsilon) \quad \text{as } \epsilon \rightarrow 0$$

*holds for  $p(\mu) \leq j \leq r(\mu)$ .*

*Proof* is analogous to that of Theorem 1. ■

Let us mention in conclusion that the results derived here raise some interesting questions, for instance, what is the following term in the expansion, what the asymptotic behaviour looks like for non-smooth surfaces, and whether similar formulae are valid in the case of  $\text{codim } \Sigma = 2, 3$  when the corresponding generalized Schrödinger operator has to be defined by means of appropriate boundary conditions.

## Acknowledgments

The authors are grateful for the hospitality extended to them, P.E. in the Department of Mathematics, Tokyo Metropolitan University, and K.Y. in the Nuclear Physics Institute, AS CR; during these visits a part of this work was done. The research has been partially supported by GAAS and the Czech Ministry of Education within the projects A1048101 and ME482. Useful comments by the referees are also appreciated.

## References

- [1] S. Agmon: *Lectures on Elliptic Boundary Value Problems*, Van Nostrand, Princeton 1965.
- [2] S. Albeverio, F. Gesztesy, R. Høegh-Krohn, H. Holden: *Solvable Models in Quantum Mechanics*, Springer, Heidelberg 1988.
- [3] S. Albeverio, P. Kurasov: *Singular Perturbations of Differential Operators*, London Mathematical Society Lecture Note Series 271, Cambridge Univ. Press 1999.

- [4] J.F. Brasche, A. Teta: Spectral analysis and scattering theory for Schrödinger operators with an interaction supported by a regular curve, in *Ideas and Methods in Quantum and Statistical Physics*, Cambridge Univ. Press 1992; pp. 197-211.
- [5] J.F. Brasche, P. Exner, Yu.A. Kuperin, P. Šeba: Schrödinger operators with singular interactions, *J. Math. Anal. Appl.* **184** (1994), 112-139.
- [6] P. Exner: Spectral properties of Schrödinger operators with a strongly attractive  $\delta$  interaction supported by a surface, in *Proceedings of the NSF Summer Research Conference (Mt. Holyoke 2002)*; AMS “Contemporary Mathematics” Series, Providence, R.I., 2003; to appear
- [7] P. Exner, T. Ichinose: Geometrically induced spectrum in curved leaky wires, *J. Phys.* **A34** (2001), 1439-1450.
- [8] P. Exner, S. Kondej: Curvature-induced bound states for a  $\delta$  interaction supported by a curve in  $\mathbb{R}^3$ , *Ann. H. Poincaré* **3** (2002), 967-981.
- [9] P. Exner, S. Kondej: Bound states due to a strong  $\delta$  interaction supported by a curved surface, *J. Phys.* **A36** (2003), 443-457.
- [10] P. Exner, K. Yoshitomi: Band gap of the Schrödinger operator with a strong  $\delta$ -interaction on a periodic curve, *Ann. Henri Poincaré* **2** (2001), 1139-1158.
- [11] P. Exner, K. Yoshitomi: Asymptotics of eigenvalues of the Schrödinger operator with a strong  $\delta$ -interaction on a loop, *J. Geom. Phys.* **41** (2002), 344-358.
- [12] P. Exner, K. Yoshitomi: Persistent currents for the 2D Schrödinger operator with a strong  $\delta$ -interaction on a loop, *J. Phys.* **A35** (2002), 3479-3487.
- [13] A. Il’in: *Matching of Asymptotic Expansions of Solutions of Boundary Value Problems*, Translations of Mathematical Monographs, Vol. 102, American Mathematical Society, Providence, R.I., 1992.
- [14] T. Kato: *Perturbation Theory for Linear Operators*, 2nd edition, Springer, Heidelberg 1976.
- [15] J.L. Lions, E. Magenes: *Non-Homogeneous Boundary Value Problems and Applications*, Vol. I, Springer, Heidelberg 1972.